AUTHOR INDEX

Arthur, Don R., recent developments in tick control in Britain, 321–30.
Asker, A. A., *see* Ragab, M. T.
Bailey, G. L., fodder-beet. III. Feeding trials with cattle, 42–48.
Barclay, C., self-cleaning, miniature thresher, 190–2.
Barrowman, J. R., castration of lambs: comparison of techniques, 193–203.
Beckley, V. R. S., *see* Pereira, H. C.
Boaz, T. G., *see* Barrowman, J. R.
Bolton, J. L., breeding alfalfa in Canada, 241–54.
Bonsma, J. C., climatological research on animal husbandry, 154–75.
Burnett, G. S., *see* Robinson, K. L.
Carey, T. M., manuring of sugar-cane, 99–115.
Castle, M. E., *see* Bailey, G. L.
Clinton, P. K. S., *see* Gibson, I. A. S.
Coey, W. E., *see* Robinson, K. L.
Constable, D. H., manuring of *Hevea* in Ceylon, 131–6.
Cormack, M. W., *see* Bolton, J. L.
Davidson, T. M. W., clonal variation and yield in potatoes, 137–40.
Dowson, W. J., *see* Wiehe, P. O.
Foot, A. S., *see* Bailey, G. L.
Gibson, I. A. S., pre-emergence seed-bed losses in groundnuts, 226–35.
Hafez, E. S. E., conception-rate and periodicity in the buffalo, 15–21.
Hafez, E. S. E., puberty in female farm animals, 217–22.
Hattingh, E. R., observations on ecology of *Steobe vulgaris*, 309–13.
Hauser, Georg F., sugar-beet cultivation in Afghanistan, 283–8.
Haylett, D. G., *see* Theron, J. J.
Hodnett, G. E., sugar-cane responses to fertilizers in Trinidad, 204–16.
Hodnett, G. E., *see* Constable, D. H.
Hofmeyr, J. H., *see* Bonsma, J. C.
Jones, K. Wilson, witchweed control. I. Hormone weed-killers, 331–9.
Kadi, M. R., *see* Ragab, H. T.
Koncar, L., wool fineness of lambs, 61–64.
Lawley, D. N., *see* Davidson, T. M. W.
MacLean, J. A. R., chemical aspects of 'black-pod' cacao disease, 340–9.
Mahedevan, P., production statistics of Sinhala cattle, 55–60.
Marle, J. van, *see* Bonsma, J. C.
Milton, W. E. J., herbage palatability on undeveloped grasslands, 116–22.
Motta, M. S., *Panicum maximum*, 33–41.
Mundy, H. G., arable farming problems in Southern Rhodesia, 123–30.
Nye, P. H., value of fertilizers in the Gold Coast. I. Methods, 176–183. II. Soils of the far north, 262–74.
Pande, H. K., effects of weeding on wheat yield, 297–303.
Pereira, H. C., grass establishment on eroded soil, 1–14.
Pugsley, A. T., back-crossing wheat for bunt resistance, 314–20.
Ragab, M. T., factors affecting weights of Ossimi lambs, 304–8.
Robinson, K. L., responses of pigs to dietary penicillin, 275–82.
Robinson, P., *see* Carey, T. M.
Russell, T. A., spacing of Nigerian cacao, 145–53.
Sheffield, F. M. L., virus diseases of sweet potato in Africa, 184–9.
Shepherd, R. W., mechanical thinning of sugar-beet, 236–40.
Smalcelj, I., *see* Koncar, L.
Soper, M. H. R., farm practice and the productivity of field beans, 22–32.
Spedding, C. R. W., effect of sub-clinical worm-burden on lambs, 255–61.
Speyer, F. C. O., reflections on the world nitrogen position, 49–54.
Theron, J. J., regeneration of humus under a grass ley, 86–98.
Thomson, L. B., the P.F.R.A. and a new prairie agriculture, 289–96.
Towers, K. G., *see* Barrowman, J. R.
Vine, H., maintenance of soil fertility in Nigeria, 65–85.
Wiehe, P. O., bacterial disease of cassava in Nyasaland, 141–3.
Wilson Jones, K., *see* Jones, K. Wilson.

SUBJECT INDEX

Agriculture, prairie, the P.F.R.A. and new, 289–96; conservation of soil and water resources in Alberta and Saskatchewan, 289–95; irrigation, major projects, 293–5; pastures, community, on abandoned land, 292–3; permanent agriculture developed, 289–93; Prairie Farm Rehabilitation Act (P.F.R.A.) and agricultural improvement in western Canada, 289 sqq.; soil drifting and soil moisture, 289–1; special projects in British Columbia and Manitoba, 294–5; strip cropping, 290; water development, 291.

Alfalfa, breeding in Canada, 241–54; cold resistance, 242; disease resistance, 243–4, 246–7; drought tolerance, 244; isolation plots and cages, 247–8; organization of projects, 241–2; selection methods, 245–6; special screening and breeding techniques, 246–52; synthetic combinations based on progeny performance, 251–2; winter hardiness, 242.

Animal husbandry, climatological research on: beef-cattle in South Africa, 154–75; adaptability in different breeds, 157 sqq.; adjustment to semi-arid sub-tropical

AUTHOR INDEX

Arthur, Don R., recent developments in tick control in Britain, 321–30.
Asker, A. A., *see* Ragab, M. T.
Bailey, G. L., fodder-beet. III. Feeding trials with cattle, 42–48.
Barclay, C., self-cleaning, miniature thresher, 190–2.
Barrowman, J. R., castration of lambs: comparison of techniques, 193–203.
Beckley, V. R. S., *see* Pereira, H. C.
Boaz, T. G., *see* Barrowman, J. R.
Bolton, J. L., breeding alfalfa in Canada, 241–54.
Bonsma, J. C., climatological research on animal husbandry, 154–75.
Burnett, G. S., *see* Robinson, K. L.
Carey, T. M., manuring of sugar-cane, 99–115.
Castle, M. E., *see* Bailey, G. L.
Clinton, P. K. S., *see* Gibson, I. A. S.
Coey, W. E., *see* Robinson, K. L.
Constable, D. H., manuring of *Hevea* in Ceylon, 131–6.
Cormack, M. W., *see* Bolton, J. L.
Davidson, T. M. W., clonal variation and yield in potatoes, 137–40.
Dowson, W. J., *see* Wiehe, P. O.
Foot, A. S., *see* Bailey, G. L.
Gibson, I. A. S., pre-emergence seed-bed losses in groundnuts, 226–35.
Hafez, E. S. E., conception-rate and periodicity in the buffalo, 15–21.
Hafez, E. S. E., puberty in female farm animals, 217–22.
Hattingh, E. R., observations on ecology of *Steobe vulgaris*, 309–13.
Hauser, Georg F., sugar-beet cultivation in Afghanistan, 283–8.
Haylett, D. G., *see* Theron, J. J.
Hodnett, G. E., sugar-cane responses to fertilizers in Trinidad, 204–16.
Hodnett, G. E., *see* Constable, D. H.
Hofmeyr, J. H., *see* Bonsma, J. C.
Jones, K. Wilson, witchweed control. I. Hormone weed-killers, 331–9.
Kadi, M. R., *see* Ragab, H. T.
Koncar, L., wool fineness of lambs, 61–64.
Lawley, D. N., *see* Davidson, T. M. W.
MacLean, J. A. R., chemical aspects of 'black-pod' cacao disease, 340–9.
Mahedevan, P., production statistics of Sinhala cattle, 55–60.
Marle, J. van, *see* Bonsma, J. C.
Milton, W. E. J., herbage palatability on undeveloped grasslands, 116–22.
Motta, M. S., *Panicum maximum*, 33–41.
Mundy, H. G., arable farming problems in Southern Rhodesia, 123–30.
Nye, P. H., value of fertilizers in the Gold Coast. I. Methods, 176–183. II. Soils of the far north, 262–74.
Pande, H. K., effects of weeding on wheat yield, 297–303.
Pereira, H. C., grass establishment on eroded soil, 1–14.
Pugsley, A. T., back-crossing wheat for bunt resistance, 314–20.
Ragab, M. T., factors affecting weights of Ossimi lambs, 304–8.
Robinson, K. L., responses of pigs to dietary penicillin, 275–82.
Robinson, P., *see* Carey, T. M.
Russell, T. A., spacing of Nigerian cacao, 145–53.
Sheffield, F. M. L., virus diseases of sweet potato in Africa, 184–9.
Shepherd, R. W., mechanical thinning of sugar-beet, 236–40.
Smalcelj, I., *see* Koncar, L.
Soper, M. H. R., farm practice and the productivity of field beans, 22–32.
Spedding, C. R. W., effect of sub-clinical worm-burden on lambs, 255–61.
Speyer, F. C. O., reflections on the world nitrogen position, 49–54.
Theron, J. J., regeneration of humus under a grass ley, 86–98.
Thomson, L. B., the P.F.R.A. and a new prairie agriculture, 289–96.
Towers, K. G., *see* Barrowman, J. R.
Vine, H., maintenance of soil fertility in Nigeria, 65–85.
Wiehe, P. O., bacterial disease of cassava in Nyasaland, 141–3.
Wilson Jones, K., *see* Jones, K. Wilson.

SUBJECT INDEX

Agriculture, prairie, the P.F.R.A. and new, 289–96; conservation of soil and water resources in Alberta and Saskatchewan, 289–95; irrigation, major projects, 293–5; pastures, community, on abandoned land, 292–3; permanent agriculture developed, 289–93; Prairie Farm Rehabilitation Act (P.F.R.A.) and agricultural improvement in western Canada, 289 sqq.; soil drifting and soil moisture, 289–1; special projects in British Columbia and Manitoba, 294–5; strip cropping, 290; water development, 291.

Alfalfa, breeding in Canada, 241–54; cold resistance, 242; disease resistance, 243–4, 246–7; drought tolerance, 244; isolation plots and cages, 247–8; organization of projects, 241–2; selection methods, 245–6; special screening and breeding techniques, 246–52; synthetic combinations based on progeny performance, 251–2; winter hardiness, 242.

Animal husbandry, climatological research on: beef-cattle in South Africa, 154–75; adaptability in different breeds, 157 sqq.; adjustment to semi-arid sub-tropical

environment, 167–70; body-temperature reaction and heat tolerance, 161–3; characteristics of cattle suited to dry, warm conditions, 171–4; comparison of Africander, British, and cross-bred cattle, 157 sqq.; fertility, 159–64; mortality rates, 165–8; weight changes, 160–3.

Animals, farm, puberty in female, 217–25; age and weight at puberty, 218–22; sheep, rabbits, and poultry in Egypt, puberty in, 218 sqq.; species, breed and individual differences observed, 218–24.

Beans, field, influence of farm practice on productivity, 22–32; bird damage, 29–30; fertilizing and manuring, 24–26; pests and diseases, 29–30; rate and method of sowing, 27–28; seed quality, 28–29; sowing date, 26–27.

Beef-cattle production in colonial territories: *see* Animal husbandry, climatological research on.

Buffalo, Egyptian dairy, conception-rate and periodicity in, 15–21; breeding efficiency and conception-rate, 16–17, 18–20, oestrous cycle length, 17–18; silent heat, 17, 20.

Bunt, resistance of wheat to, *see* Wheat, back-crossing of.

Cacao, Amelonado, some chemical aspects of 'black-pod' disease in, 340–9; alkaloids (theobromine, &c.) and nitrogen content of healthy and diseased beans, 342–3, 346; changes in composition with time, 343–4; fat and free fatty acid content, 341–3, 346; incidence of 'black-pod' disease in Gold Coast and Nigeria, 343–5; sugars in healthy and diseased beans, 343, 347; utilization of diseased beans, 347–8.

Cacao, Nigerian, spacing of, 145–53; farmers' planting method, 151; peasant farms, 149–51; trial begun in 1934, 145–7; trial begun in 1942, 148–9; value of close planting and closed canopy, 152–3.

Cassava, a bacterial disease of, in Nyasaland, 141–3; description of pathogen (*Xanthomonas cassava* sp. nov.), 142–3; leaf-spot and defoliation resulting from, 141.

Cattle, beef-; *see* Animal husbandry.

Cattle, Sinhala, of Ceylon: general life and production statistics, 55–60; breeding, 56–57; calving, lactation and dry-period statistics, 56; heritability of milk yield, 59; milk production 57–59.

Dura (sorghum), witchweed in, *see* Witchweed (*Striga hermontheca*), control of.

Farming, arable, in Southern Rhodesia, some problems of, 123–30; compost, 126; dung, value of, 125, 127; grass leys and legumes, 128–9; green manuring, 125 sqq.; nitrogen and phosphate problems, 129–30; soil fertility maintenance, 125; soil organic matter, 126–7.

Fertilizers, survey of the value of, in Gold Coast food-farming areas. I. Aims and methods, 176–183; conditions in the areas, 176–9; field experiment methods used, 178–82; organization of experiments, 182. II. Granitic soils of the far north, 262–74; cash return from fertilizers, 270–2; cereals and groundnuts, results with, 264 sqq.; climate, soils, and agriculture; 262–4; fertilizer recommendations, 272–3; nitrogen and phosphate, responses to, 264 sqq.; organic manuring, effects of, 266–7.

Fodder-beet. III. Feeding trials with dairy cattle, 42–48; composition of fodder-beet and other feeds, 44; cows, dairy, direct feeding trials, 42–43; mangolds, comparison with fodder-beet, 43–48.

Grass establishment on eroded soil in Kenya, 1–14; carrying capacity of experimental pastures, 7–9; conservation of soil moisture, 10–11; establishment of grasses with soil-conserving cultivation methods, 5–8; labour expenditure, 12–13; overgrazing, and potential carrying capacity, 2–3; soil erosion, severity of, 1–2; soil restoration methods used, 3–4; soil structure studies, 11–12.

Grass ley, *see* Soil humus, regeneration of.

Grasslands, undeveloped, in Wales: palatability of herbage, 116–22; grasses, herbs and other species, discussion of, 120–1; palatability, relative, of species on open hill grazings, 117–20; season, influence of, 117–21.

Groundnuts, pre-emergence seed-bed losses of, in Tanganyika, 226–35; fungi as a cause of pre-emergence loss, 229–32; mechanical injury to seed, 226–9; Rhizopus spp., chief cause of loss, 230–2; seed-treatment with fungicides, effect of, 232-5.

Guinea grass, *see Panicum maximum*.

Hevea brasiliensis, manuring of, in Ceylon, 131–6; girth of trees, fertilizer effects on, 132; nitrogen, phosphate and potash responses and interactions, 132–5; phosphate, importance of, 132–4; yield of rubber, fertilizer effects on, 133 sqq.

Lambs, castration of, comparison of techniques, 193–203; crushing (the Burdizzo instrument), 195; physical effects on lambs, 196–7; rubber-ring ligature (Elastrator), 195–6; subsequent performance of lambs, 197–9.

Lambs, Ossimi, factors affecting weights of, 304–8; genetic correlations between weights, 306–7; heritability of birth, weaning, and marketing weights, 305 sqq.; phenotypic correlations, 305–6.

Lambs, sub-clinical worm-burden of, effect on liveweight gain, 255–61; faeces consistency, food intake, and egg-counts: effects of dosing with worm larvae, 259–60; *Trichostrongylus axei*, effects on liveweight changes, &c., of dosing with larvae of, 256 sqq.

Lambs, wool of, fineness and medullate fibres in, 61–64; age and region of lamb, relation of fibre thickness to, 61 sqq.; relation of medullated fibres to, 63–64; Yugoslavia, properties of wool of cross-bred lambs in, 61 sqq.

Lucerne, *see* Alfalfa, breeding in Canada.

Nitrogen, reflections on the world position

of, 49–54; future trends in demand for agricultural nitrogen, 50 sqq.; population and cropped area of world, in relation to potential nitrogen consumption, 52 sqq.; potential nitrogen in A.D. 2000, 53–54; present production and consumption of fertilizer nitrogen, 49–50.

Panicum maximum, 33–41; distribution of Guinea grass in tropical countries, 33; establishment, 35; management, utilization and conservation, 35–37; nutritional status, 39–40; strains and ecotypes, 32–34; yields and responses to fertilizers, 37–39.

Pigs, response of, to dietary procaine penicillin, 275–82; appetite, effect of penicillin on, 276, 278–9; feeding trials with weanling and fattening pigs, 276–8; food utilization, 277, 279; maximum response, time of, 276–81; recession due to discontinuance of penicillin, 277; weight-gains, effect of penicillin on, 277 sqq.

Potato, sweet, virus diseases of, in Africa, 184–9; distribution in tropical Africa, 184–5, 187–9; other conditions resembling virus disease, 186; spread of virus, 187–9; symptoms of virus disease in sweet potato, 185–6.

Potatoes, evidence of clonal variation affecting yield of, 137–40; clones of King Edward variety, yields of, 138–9; yield differences between clones, significant and heritable, 138 sqq.

Rubber, manuring of, *see Hevea brasiliensis*.

Soil fertility, experiments on the maintenance of, in Nigeria, 65–85; burning mucuna (green manure), effect on fertility, 73–83; fallows, vegetative, restoration by, 83, 84; fertilizer (nitrogen and phosphate) experiments and results, 76–79; green manure, use of in fertility maintenance, 68 sqq.; mucuna (velvet bean) as main green manure crop, 69–84; soil analyses, 81–83.

Soil humus, regeneration of, under a grass ley, 86–91; fertilizer (nitrogen and phosphate) responses, as hay yields, 87–88; hay and protein production in relation to soil humus, 91–96; nitrogenous fertilizers, use of to increase humus formation, 91–97; Rhodes grass ley in South Africa, experiment with, 86 sqq.

Stoebe vulgaris, observations on the ecology of, 309–13; control of Stoebe by heavy grazing with increased fertility, 311 sqq.; distribution of *Stoebe vulgaris* (*Slangbos*) in S. Africa, 329–11; management, effects of, 311–12; mowing or burning relatively ineffective, 311; role of Stoebe in succession, 311; soil fertility, effect of, 312–3.

Striga hermontheca, *see* Witchweed.

Sugar-beet cultivation and improvement in Afghanistan, 283–8; comparison of native and improved cultivation methods, 285–7; native cultivation method described, 284; sowing in rows on flat soil, success of, 286 sqq.

Sugar-beet, mechanical thinning of, 236–40; conditions for success with mechanical thinning, 238 sqq.; cross-blocking, trials with, 236–8; precision drills, 237–8; rubbed seed, 237–8.

Sugar-cane, manuring of, 99–115; Commonwealth territories included in survey, 99; designs and numbers of field experiments used, 99; fertilizer (nitrogen, phosphate, and potash) effects on cane yield, 105–11; liming responses, 114–15; nitrogen responses, 105–8; organic manures (pen and green manures, molasses), 111–14; response curves and optimal dressings, 101–5; sugar content, fertilizer effects on, 100–1.

Sugar-cane, responses to fertilizers in Trinidad, 204–16; designs of experiments and conversion of results, 206–9; molasses and filter-press mud, effects of, 213–14; nitrogen responses, 209–10; pen manure, effects of, 212–13; phosphate responses 210–11; potash responses, 211–12; sugar content, fertilizer effects on, 213–14.

Sweet potatoes, *see* Potato, sweet.

Thresher, miniature, a self-cleaning, 190–2; corn and peas, threshing single ears and pods, 190 sqq.

Tick control in Britain, recent developments in, 321–30; arsenic for tick control, 323–5; cattle ticks, control of, 322 sqq.; chemical treatment of livestock, 323 sqq.; chlorinated hydrocarbons (BHC, DDT, &c.), 327–9; cultural practices for tick control, 321–2; derris and pyrethrum, 325–7; sheep ticks, control of, 321 sqq.

Wheat, back-crossing of, for resistance to bunt in South Australia, 314–20; Doubbi tetraploid wheat, resistance and incorporation of, 314–15; genes in wheat resistant to 'races' of bunt, 319–20; Insignia series of crosses, 315–17; Javelin series, 317–18; Ranee-Doubbi selections as basis of resistance, 314 sqq.; Ridley series of crosses, 318–19.

Wheat, weeding, in India, effect on the yield of, 297–303; density, relative, of weeds and wheat, 298; fertility of soil, interaction with effect of weeding, 302–3; hand-hoe and hand-pulling, comparison of, 299 sqq.; tillering, effect of weeding on, 299 sqq.; weed species, names and density of, 298; yield of grain, effect of weeding on, 298 sqq.

Witchweed (*Striga hermontheca*), control of by hormone weed-killers, 331–9; 2,4-D for witchweed control in Sudan, design and results of experiments, 331 sqq.; action of 2,4-D on Striga, 336–8; damage to dura, extent of and recovery from, 332–3; dura (sorghum), irrigated, control of witchweed in, 331 sqq.; dura varieties, relation to effectiveness of weed-killer, 333–8; strength of weed-killer, effects on dura and witchweed, 332–7; Striga observations and weights, 333–5; time of application, importance of, 333–8.

Wool, fineness of, *see* Lambs, wool of.

Worm-burden, sub-clinical, *see* Lambs, sub-clinical worm-burden of.